SEXUAL PRACTICES, THEIR INFLUENCERS AND UTILIZATION OF HIV
SERVICES AMONG FEMALE SEX WORKERS IN MOMBASA COUNTY, KENYA

ROBERT ABUGA MOKINU

A THESIS SUBMITTED TO THE SCHOOL OF APPLIED AND HEALTH SCIENCES IN THE DEPARTMENT OF ENVIRONMENTAL AND HEALTH SCIENCES IN PARTIAL FULFILLMENT OF THE REQUIREMENTS FOR THE AWARD OF MASTER DEGREE IN PUBLIC HEALTH OF TECHNICAL UNIVERSITY OF MOMBASA

# **DECLARATION**

This thesis is my original work and has not been presented for a degree award in any other university.

Robert Abuga Mokinu	
MPH/0025/2019	
Signature	Date
This thesis has been presented with our appro	oval as university supervisors.
Dr. Shadrack Yonge, PhD	
Signature	Date
Prof. Peter Gichangi, Ph.D.	
1101. Teter Gichangi, Th.D.	
Signature	Date

## **DEDICATION**

I dedicate this study to my daughter Sharon Kerubo Abuga, my mother Yovensia Nyaboke Mokinu and my late father Nahashon Mokinu. Daddy's encouragement "Kari motwe" (keep moving on) has taken me this high, his legacy lives.

### ACKNOWLEDGMENT

I gratefully acknowledge all of the faculty members at Technical University of Mombasa department of Environmental and Health Sciences for their invaluable assistance. Dr. Shadrack Yonge and Prof. Peter Gichangi deserve special mention for their professional assistance and guidance. Your advice and technical assistance have been invaluable and inspirational to me. Finally, I would like to express my gratitude to the International Centre for Reproductive Health – Kenya (ICRH-K) for their support (accepted this study to use part of their data).

## TABLE OF CONTENTS

DECLARATIONii
DEDICATIONiii
ACKNOWLEDGMENTiv
TABLE OF CONTENTSv
LIST OF TABLESx
LIST OF FIGURESxi
LIST OF ABBREVIATIONS AND ACRONYMSxii
DEFINATION OF TERMSxiv
ABSTRACTxv
CHAPTER ONE1
INTRODUCTION1
1.1 Background Information of the Study1
1.2 Statement of the Problem
1.3 Justification4
1.4 Research Objectives5
1.4.1 General Objective5
1.4.2 Specific Objectives5
1.5 Null Hypotheses5
1.6 Significance of the Study6
1.7 Study Scope6
1.8 Study Limitations6

1.9 Conceptual Framework	7
CHAPTER TWO	8
LITERATURE REVIEW	8
2.1 Global Overview	8
2.2 Socio-Demographic Characteristics of Female Sex Workers in Mombas	a9
2.2.1 Young Age and Sex Work	9
22.2 Education Level of Female Sex Workers	10
2.2.3 Marital Status of Female Sex Workers	11
2.2.4 Residence of Female Sex Workers	12
2.2.5 Income	12
2.2.6 Peer Pressure Influence	13
2.3 Sexual Practices Among Female Sex Workers	14
2.3.1 Anal Intercourse	14
2.3.2 Oral Sex	15
2.3.3 Condoms Use and Non-Use during Sex	16
2.3.4 Multiple Sexual Partners	17
2.3.5 Forced Sex Among Female Sex Workers	17
2.4 Alcohol Use by Female Sex Workers	18
2.5 Violence Against Female Sex Workers	19
2.5.1 Physical Violence	19
2.5.2 Sexual Violence	20
2.5.3 Psychological Violence	20
2.6 Utilization of HIV Services	20

2.6.1 HIV Testing	21
2.6.2 Antiretral Viral Therapy (ART) Accessibility and Usage	22
2.6.3 Pre-Exposure Prophylaxis (Prep) Use	23
2.6.4 Use of Post-Exposure Prophylaxis (PEP)	24
CHAPTER THREE	26
MATERIALS AND METHODS	26
3.1 Data Source	26
3.2 Study Location	26
3.3 Design of the Study	26
3.4 Population of the Study	27
3.4.1 Inclusion Criteria	27
3.4.2 Criteria for Exclusion	27
3.4.3 Independent Variables	28
3.4.4 Intervening Variables	28
3.4.5 Dependent Variable	28
3.5 Sample Size	28
3.6 Sampling Technique	29
3.7 Tools for Data Collection	29
3.7.1 Pre-Testing of Research Instruments	30
3.7.2 Validity	30
3.7.3 Reliability	30
3.8 Procedure for Data Collection	30
3.9 Management and Analysis of Data	31

3.10 Ethical Considerations
CHAPTER FOUR
RESULTS AND DISCUSSION
4.1 Socio-Demographic Characteristics of FSWs in Mombasa County Between 2014 - 2015
4.2 Sexual Practices
4.2.1 Female Sex Workers' Behavior in Mombasa County35
4.2.2 Frequency and Type of Sexual Intercourse with Partners35
4.2.3 Type of Condoms used by Female Sex Workers in Mombasa County between 2014 - 2015
4.3 Influencers of Sexual Practices Among Female Sex Workers
4.3.1 Alcohol use and Violence Among Female Sex Workers
4.4 Utilization of HIV Services by FSWs in Mombasa County
4.4.1 Knowledge of HIV Prevention and Testing by FSWs40
4.4.2 Attitudes Toward HIV Testing Among FSWs
4.5 Determinants of Risky Sex Practice among FSWs in Mombasa County4.5.1 Association of FSWs Socio-Demographic Characteristics, Behaviors and
Frequency of Sex or Sex Type with Risky Sex Practices
4.5.2 Alcohol use and Violence Factors Associated with Risky Sex Practice 49
4.5.3 Knowledge HIV Prevention and Testing Factors Associated with Risky Sex Practice
4.5.4 Attitudes towards HIV Testing Factors Associated with Risky Sex Practices 54
4.6 Predictors of Risky Sex Practice among FSWs in Mombasa County54
4.7 Discussion

4.7.1 Introduction	7
4.7.2 Socio-Demographic Characteristics of FSWs	7
4.7.3 Sexual Practices Among Female Sex Workers	7
4.7.4 Influencers of Sexual Practices among Female Sex Workers	)
4.7.5 Utilization of HIV Services	<u>)</u>
CHAPTER FIVE64	Ł
CONCLUSION AND RECOMMENDATIONS64	Ł
5.1 Conclusion	Į
5.2 Recommendations65	5
REFERENCES	ĵ
APPENDICES88	3
Appendix I: Request and approval for secondary data from boresha project 88	3
Appendix II: Map of study area	)
Appendix III: Questionnaire90	)

# LIST OF TABLES

Table 4.1: Socio-demographic characteristics of FSWs in Mombasa County between
2014 - 2015
Table 4. 2: Female sex workers' behaviour, frequency and type of sexual intercourse
with partners in Mombasa County between 2014 - 201536
Table 4.3: Alcohol use and violence among female sex workers in Mombasa County
between 2014 - 2015
Table 4.4: Knowledge of HIV prevention and testing by FSWs in Mombasa County
between 2014 - 2015
Table 4.5: Attitudes toward HIV testing among FSWs in Mombasa County between
2014 - 2015
Table 4.6 (b): Association of FSWs Socio-Demographic Characteristics, Behaviors and
Frequency of Sex / Sex Type with Risky Sex Practices in Mombasa County between
2014 - 2015
Table 4.6 (b): Association of FSWs socio-demographic characteristics, behaviors and
frequency of sex / sex type with risky sex practices in Mombasa County between 2014
- 2015
Table 4.7: Alcohol use and violence factors associated with risky sex practice in
Mombasa County between 2014 - 2015 51
Table 4.8: Knowledge of HIV prevention and testing factors associated with risky sex
practice in Mombasa County between 2014 - 2015 53
Table 4.9: Predictors of risky sex practice among FSWs in Mombasa County in 201556

### LIST OF FIGURES

Figure 1.1: Conceptual	framework	k

LIST OF ABBREVIATIONS AND ACRONYMS

AI Anal Intercourse

AIDS Acquired Immunodeficiency Syndrome

**ART** Antiretroviral Therapy

**ARV** Antiretroviral

**ERC** Ethical Review Committee

**FSWs** Female Sex Workers

**HIV** Human Immune Deficiency Virus

**HPV** Human Papilloma Virus

ICRH International Center for Reproductive Health

**IP** Intimate Partner

**KNBS** Kenya National Bureau of Statistics

**KNH** Kenyatta National Hospital

**KP** Key Population

MARPs Most at Risk Populations

**MPH** Masters in Public Health

**MSM** Men who have Sex with Men

NACOSTI National Commission for Science, Technology and Innovation

NASCOP National AIDS and STI Control Program

**PLHIV** People Living with HIV

PrEP Pre-Exposure ProphylaxisPWID People Who Inject Drugs

**REF** Risky Sexual Behaviour

SD Standard Deviation

**SRH** Sexual and Reproductive Health

STIs Sexually Transmitted Infections

STDs Sexually Transmitted Diseases

TLD Tenofovir Lamivudine Dolutegravir
TUM Technical University of Mombasa

**UNAIDS** United Nations Program on HIV/AIDS

VI Vaginal Intercourse

WHO World Health Organization

### **DEFINATION OF TERMS**

Early sexual debut Having first incident of sexual under the age of 15 years

Risky sex

The activity that will increase the probability that a person engaging in sexual activity with another person infected with a sexually transmitted infection will be infected

Sex work

The exchange of sexual services, performances, or products for material compensation.

### **Prostitution**

The practice of engaging in relatively indiscriminate sexual activity, in general with someone who is not a spouse or a friend, in exchange for immediate payment in money or other valuables

Female sex workers

Girls and women who regularly or occasionally receive money in exchange for sexual services.

#### **ABSTRACT**

Female sex workers (FSWs) are vulnerable to acquiring HIV and other sexually transmitted infections due to risky sexual practices. Risky sexual practices refer to sexual acts and behaviours that increase the chances of contracting and transmitting sexually transmitted infections. Understanding and addressing the gaps in safer sexual practices among FSWs can help to reduce HIV acquisition and transmission. This study focused on the sexual practices, their determinants, HIV knowledge, and use of HIV testing services among FSWs in Mombasa County, Kenya. Cross-sectional research design was adopted in the investigation. Respondents were recruited by systematic random selection at predetermined intervals from bars and clubs until a sample size of 160 was reached. Women who were above 18 years and had in the past exchanged goods or money for sex were referred to as female sex workers. Descriptive statistics and inferential analysis using R was done and p<0.05 was regarded statistically significant. Ethical approval and clearance were obtained from the Kenyatta National Hospital and University of Nairobi Ethics Review Committee. Informed consent was obtained from participants prior to the interviews. Most of the participants (98.74%) were unmarried, 97.48% reported vaginal intercourse, 50.96% reported using alcohol before sex and 10.69% had a tertiary education. Risky sex was practiced by 27.67% of study participants. Slightly more than a half (64.15%) had tested for HIV within three months, 13.84% believed that it is safe to reuse condoms and 10.06% that it is safe engaging in unprotected sex. Risky sexual practices in bi-variate analysis were associated with frequency of sexual intercourse, frequency of vaginal intercourse, alcohol use, violence and poor HIV knowledge. In multivariate analysis, risky sexual practices were associated with frequency of sexual intercourse, alcohol use and poor HIV knowledge. Female sex workers are still engaging in unprotected sex while intoxicated, belief in re-using condoms and high frequency of sexual intercourse. Alcohol use significantly correlates to risky sexual practices. Few FSWs test for HIV within three months and some still has poor knowledge about HIV. Interventions to address these modifiable factors such as promoting partners' conversations about condom use, routine alcohol use risk reduction counselling and creating FSWs social networks are needed.