

Type IVB Pilus Operon Promoter Controlling Expression of the Severe Acute Respiratory Syndrome-Associated Coronavirus Nucleocapsid Gene in *Salmonella enterica* Serovar Typhi Elicits Full Immune Response by Intranasal Vaccination[∇]

Fengling Luo,^{1†} Yong Feng,^{1,2†} Min Liu,¹ Pingfei Li,¹ Qin Pan,¹ Victor Tunje Jeza,¹ Bing Xia,³ Jianguo Wu,^{2*} and Xiao-Lian Zhang^{1*}

Department of Immunology, Hubei Province Key Laboratory of Allergy and Immune-Related Diseases, The State Key Laboratory of Virology, Wuhan University School of Medicine, Wuhan 430071, People's Republic of China¹; The State Key Laboratory of Virology, College of Life Sciences, Wuhan University, Wuhan 430072, People's Republic of China²; and Zhongnan Hospital, Wuhan University, Wuhan 430071, People's Republic of China³

Received 10 February 2007/Returned for modification 6 March 2007/Accepted 31 May 2007

Attenuated *Salmonella enterica* serovar Typhi strains have been considered to be attractive as potential live oral delivery vector vaccines because of their ability to elicit the full array of immune responses in humans. In this study, we constructed an attenuated *S. enterica* serovar Typhi strain stably expressing conserved nucleocapsid (N) protein of severe acute respiratory syndrome-associated coronavirus (SARS-CoV) by integrating the N gene into the *pilV* gene, which was under the control of the type IVB pilus operon promoter in *S. enterica* serovar Typhi. BALB/c mice were immunized with this recombinant strain through different routes: intranasally, orogastrically, intraperitoneally, and intravenously. Results showed that the intranasal route caused the highest production of specific immunoglobulin G (IgG), IgG2a, and secretory IgA, where IgG2a was imprinted as a Th1 cell bias. Moreover, this recombinant live vaccine induced significantly high levels of specific cytotoxic T-lymphocyte activities and increased gamma interferon-producing T cells compared with the parental strain. Our work provides insights into how the type IVB pilus operon promoter controlling SARS-CoV N gene expression in *Salmonella* might be attractive for a live-vector vaccine against SRAS-CoV infection, for it could induce mucosal, humoral, and cellular immune responses. Our work also indicates that the type IVB pilus operon promoter controlling foreign gene expression in *Salmonella* can elicit full immune responses by intranasal vaccination.

Severe acute respiratory syndrome-associated coronavirus (SARS-CoV) is a new emerging virus and has a high mortality rate along with a huge economic impact worldwide. It contains five major open reading frames encoding the replicase polyprotein, the spike (S) protein, the envelope (E) protein, the membrane glycoprotein (M), and the nucleocapsid (N) protein. N protein of SARS-CoV is a 46-kDa conserved protein that participates in the replication and transcription of the virus and interferes with the cell cycle of host cells (36). Amino acid sequence homology between SARS-CoV and other CoVs is low, and this is a possible cause for the difference in pathogenesis (25). It was reported that more than 94% of SARS patients were positive for N protein-specific antibodies that appeared at early stages of infection (41).

Other studies showed that the highest immune responses were generated by DNA vaccination with the N gene against

SARS-CoV in mice among constructs encoding N, M, and E proteins (18). Therefore, the N protein was chosen as the target antigen in this study.

Currently no vaccine is licensed for human SARS-CoV, although effective vaccines have been developed for other animal CoVs (33). Recently, several vaccine strategies have been examined for prevention against SARS infection. They include inactivated viruses, DNA vaccine, and recombinant viral vectors based on adenoviruses, baculoviruses, and parainfluenza viruses (2, 3, 13, 35). In addition, the surface-displayed SARS-CoV spike protein on *Lactobacillus casei* has been reported to induce neutralizing antibodies in mice (22, 23). Studies of animal CoV vaccines have demonstrated that both systemic and cell-mediated immune responses are important in preventing the viral infections (17). DNA vaccines targeting the N protein of SARS-CoV generated strong N-specific humoral and cellular immunity and reduced the titers of challenging vaccinia virus expressing the N protein of the SARS virus (20). SARS-CoV N protein and S protein could induce a long persistence of memory T-cell response in humans (31, 42). These pieces of evidence indicate that both humoral and cellular immune responses were vital to defense against acute SARS infection (34).

The initial SARS-CoV infection occurs primarily in the mucosal epithelial cells in the respiratory tract. There is evidence that mucosal immunity is important in the rapid immune re-

* Corresponding author. Mailing address for X.-L. Zhang: Department of Immunology, Wuhan University School of Medicine, 165 Donghu Road, Wuhan 430071, People's Republic of China. Phone: 86-27-87331183. Fax: 86-27-87336380. E-mail: ZhangXL65@whu.edu.cn. Mailing address for J. Wu: The State Key Laboratory of Virology, College of Life Sciences, Wuhan University, Wuhan 430072, People's Republic of China. Phone: 86-27-68754979. Fax: 86-27-68754592. E-mail: wu9988@vip.sina.com.

† F.L. and Y.F. contributed equally to this work.

∇ Published ahead of print on 27 June 2007.

sponse to mucosal infections (42). Thus, in order to develop an effective vaccine for SARS-CoV, a good SARS vaccine candidate should elicit both mucosal and systemic immunities.

Attenuated *Salmonella* strains have been supposed as attractive potential live oral delivery vector vaccines because of their ability to elicit a full array of immune responses in humans (1, 8, 10, 21, 24, 27). Previously, we demonstrated that a *Salmonella enterica* serovar Typhi *pilS*::Km^r mutant strain significantly attenuated the adhesion and invasion to human intestinal or monocytic cells compared with those of the wild-type strain (28). The type IVB *pil* operon in *S. enterica* serovar Typhi contains 11 genes, *pilLMNOPQRSTU*V, of which *pilS* encodes the main structural protein and *pilV* codes for pilus-tip adhesion of the type IVB pili. These genes form a transcriptional unit under the control of one *pil* promoter, as evidenced by the absence of independent promoter. In this study, we sought to obtain an effective *Salmonella*-based viral vaccine candidate with both mucosal and systemic immunities. We constructed an *S. enterica* serovar Typhi *pilS*::Km^r *pilV*::N⁺ mutant strain by integrating the N gene with the *pilV* gene under the control of *pil* operon promoter as the first step towards constructing a live oral *Salmonella*-SARS-CoV vaccine and analyzed its specific immune responses in mice with different immunization routes.

MATERIALS AND METHODS

Strains and plasmids. The attenuated *S. enterica* serovar Typhi strain Ty2 (*cys try galE-H1* via Rif^r *pilS* Kan^r) (15, 44), was the source for the vaccine strain in this study. The eukaryotic expression vector pCMV-Tag2B-N expressing full-length N gene of a SARS-CoV strain (WHU) was described previously (46).

Construction of the *S. enterica* serovar Typhi *pilS*::Km^r *pilV*::N⁺ recombinant vaccine stably expressing N protein of SARS-CoV. The most important approach used in the construction of recombinant vaccine is homologous recombination. Plasmid pUST100 containing the ApaI-EcoRV digestive site fragment with *pilV* and *rci* genes (45) was cleaved with BamHI, which generated a small fragment and a large fragment, after which the large fragment was religated to generate a new plasmid named pUST110. The N gene (1.2 kb) of SARS-CoV was obtained by digestion of pCMV-Tag2B-N with BamHI and EcoRI and then inserted into the vector pUST110 digested with BglII and EcoRI to create a recombinant plasmid, pUST110-N. Plasmid pUST110-N was transferred into the *S. enterica* serovar Typhi *pilS*::Km^r strain via the *S. enterica* serovar Typhimurium-modifying strain J357 (r⁻ m⁺) (44) with Ap^r and Km^r selection. Individual clones were then grown in 5 ml LB medium without antibiotic selection at 42°C for periods of ca. 24 h before transfer of a 0.1-ml aliquot into 5 ml fresh medium. After three or four transfers, aliquots were plated onto LB medium containing kanamycin and the resulting colonies that had lost the transformed plasmids were checked for ampicillin sensitivity. The *S. enterica* serovar Typhi *pilS*::Km^r *pilV*::N⁺ recombinant vaccine, in which the N gene was inserted into the *S. enterica* serovar Typhi genome, was screened from large numbers of colonies and examined by PCR amplification using primers adjacent to or inside the inserted fragment as well as by Western blot analysis.

PCR was used to identify the N gene in the *S. enterica* serovar Typhi *pilS*::Km^r *pilV*::N⁺ strain. The primers outside the recombinant sequences used for amplification were 5'-CGATGATAGTCCGGAATCAGC-3' and 5'-ATCCGGACG ACCATTGACCTG-3'. Primers with insert sequences used for amplification were 5'-ATGCTCTGATAATGGACC-3' and 5'-TGCCTGAGTTGAATCAG-3'. The expression of N protein in the *S. enterica* serovar Typhi *pilS*::Km^r *pilV*::N⁺ strain was determined by Western blot analysis.

Immunization process optimization. Seven- to 8-week-old female BALB/c mice were prepared for immunization. Four groups (six mice per group) of mice were immunized with the recombinant strain or the parental bacterium strain (i) intranasally (i.n.) by dispensing 10 µl of vaccine suspension containing 10⁹ CFU directly into the mouse nasal cavity; (ii) orogastrically (o.g.) by placing 100 µl of vaccine suspension containing 10⁹ CFU into the lower esophagus using a gavage needle; (iii) intraperitoneally (i.p.); and (iv) intravenously (i.v.) by injecting 100 µl of vaccine suspension containing 10⁷ CFU into the abdominal cavity and vena caudalis, respectively, on days 0 and 14. Mice were euthanized at day 14 after the

last immunization. Production of serum immunoglobulin G (IgG) against N protein was measured by enzyme-linked immunosorbent assay (ELISA).

Purification of recombinant N protein and antibody production. The N gene of SARS-CoV was generated by digestion of pCMV-Tag2B-N (46) with BamHI and EcoRI and cloned into the BamHI and EcoRI sites of prokaryotic vector pGEX-KG to yield plasmid pGEX-N. Plasmid pGEX-N was transformed into *E. coli* BL21(DE3)[pLysS]. The glutathione *S*-transferase (GST)-N fusion protein was overexpressed in *E. coli* after the induction of isopropyl-β-D-thiogalactopyranoside (IPTG). The expressed protein was purified by glutathione Sepharose 4B (Amersham Biosciences). Anti-N antibody was prepared from immunized rabbit sera with the recombinant GST-N protein.

Measurement of antibody levels by ELISA. Anti-N antibodies were detected 2 weeks after the final immunization. The levels of antibodies against N protein were determined by ELISA. Briefly, 96-well plates were coated with 100 µl of the recombinant N protein (10 µg/ml) (expressed and purified from *E. coli* BL21) in carbonate buffer, pH 9.6, for 3 h at 37°C and blocked overnight with 1% bovine serum albumin at 4°C. After incubation, the plates were washed with phosphate-buffered saline (PBS) containing 0.05% Tween 20 (PBST). Sera were subjected for titer determination in twofold serial dilutions. Antibodies bound to the immobilized antigens were detected using horseradish peroxidase-labeled anti-mouse immunoglobulin G (IgG), IgG2a (Southern Biotech) diluted at a ratio of 1:1,500 in PBST, and substrate solution containing *o*-phenylenediamine (1 mg/ml) and H₂O₂ (0.03%) in 0.1 M citrate phosphate buffer. Test and control sera were run in triplicate.

Detection of anti-N S-IgA antibody by ELISA. For detection of anti-N secretory IgA (S-IgA), briefly, flat-bottomed microtiter plates were coated with the recombinant N protein (expressed and purified from *E. coli* BL21) and blocked with 1% bovine serum albumin. The supernatant extracted from 100 µg of original fecal pellets of each mouse mixed with 100 µl of PBS was subjected to titer determination in twofold serial dilutions. Samples were incubated at 100 µl/well for 1 h at 37°C. Horseradish peroxidase-conjugated goat anti-mouse IgA antibodies (Southern Biotech) were added at a 1:5,000 dilution in PBST and incubated at 37°C for 1 h. Experimental and control samples were run in triplicate.

Stable transfection. A total of 1 × 10⁶ CT26 (*H-2^d*) murine colon tumor cells (39) were placed into a 60-mm-diameter plate 24 h before transfection. Each 60-mm-diameter plate of cells was transfected with 8 µg of plasmid pCMV-Tag2B-N by Lipofectamine 2000 (Invitrogen). For stable transfections, G418 was added to the cell culture medium at a final concentration of 0.6 mg/ml after 48 h posttransfection. Cell culture medium was changed every 2 days. After 4 weeks of selection, N-expressing CT26 cells were obtained. Cells from wells containing single clones were selected for further analysis, and individual transfected clones were tested for expression of N protein by Western blot analysis.

Specific CTL activity. Seven- to 8-week-old female BALB/c mice were immunized i.n. with 10⁹ CFU/10 µl of the recombinant strain or the parental bacterium strain on days 0 and 14, respectively. Mice were euthanized at day 28. Splenocytes from mice were resuspended in complete RPMI 1640 with 10% fetal bovine serum and analyzed for cytotoxic T-lymphocyte (CTL) activity. Twofold serial dilutions of mice splenocytes as expanded effector cells (6.25 × 10³ to 2 × 10⁵ cells/well) were stimulated by recombinant N protein (5 µg/ml, expressed and purified from *E. coli* BL21) in vitro and incubated with major histocompatibility complex (MHC)-matched CT26 (*H-2^d*), which stably expressed N protein as target cells (2 × 10⁴ cells/wells) for 4 h at 37°C in the presence of 5% CO₂. Cultures were centrifuged at 1,000 rpm for 5 min, 50 µl of supernatants per well was then transferred to enzymatic assay plates, and lysis was determined by measuring released lactate dehydrogenase (LDH) by using the Cytotoxic 96 assay kit (Promega Corp., Madison, WI). The absorbance values from the supernatants were recorded at 490 nm on an ELISA microplate reader. The data are means ± standard deviation (SD) of five different wells. The percentage of cytotoxicity was calculated as [(experimental - effector spontaneous - target spontaneous)/(target maximum - target spontaneous)] × 100, where spontaneous release is the count released by target cells in the absence of effector cells and maximal release is the number of counts released in the presence of lysis solution.

ELISPOT assay. For the enzyme-linked immunospot (ELISPOT) assay, 96-well filtration plates were coated overnight at 4°C with 50 µl (10 µg/ml) of anti-mouse gamma interferon (IFN-γ) or interleukin-4 (IL-4) in sterile PBS. The plates were blocked for 2 h at 37°C with RPMI 1640 containing 10% fetal calf serum and 1% bovine serum albumin and were washed three times with sterile PBS. Various dilutions of splenocytes from immunized or control mice in 200 µl of complete medium were placed in each well and were cultured for 18 h in RPMI 1640 alone (negative control) or with 5 µg per ml recombinant N protein (expressed and purified from *E. coli* BL21) or 8 µg per ml of concanavalin A




FIG. 1. Construction and analysis of the *S. enterica* serovar typhi *pilS::Km^r pilV::N⁺* recombinant strain with chromosomal integration of the N gene. (A) Diagram of the construction of the recombinant strain. (B) Analysis of the recombinant strain by PCR amplification using primers outside recombinant sequences. Lane 1, *S. enterica* serovar Typhi *pilS::Km^r* parental strain; lane 2, plasmid pUST110-N; lane 3, *S. enterica* serovar Typhi *pilS::Km^r pilV::N⁺* recombinant strain; lane 4, DNA marker. (C) Analysis of recombinant strain by PCR amplification using primers with insert sequences. Lane 1, DNA marker; lane 2, *S. enterica* serovar Typhi *pilS::Km^r pilV::N⁺* recombinant strain; lane 3, *S. enterica* serovar Typhi *pilS::Km^r* parental strain. (D) Examination of N protein (45 kDa) expression in the *S. enterica* serovar Typhi *pilS::Km^r pilV::N⁺* recombinant strain with anti-N antibody. Lane 1, *S. enterica* serovar Typhi *pilS::Km^r* parental strain; lane 2, *S. enterica* serovar Typhi *pilS::Km^r pilV::N⁺* recombinant strain.

(positive control) in triplicate wells incubated at 37°C for 24 h. Plates were washed with PBS containing 0.025% Tween 20 and were overlaid with 50 μ l (5 μ g/ml) of biotinylated anti-mouse IFN- γ or IL-4. The plates were washed six times with PBS containing 0.025% Tween 20 and were treated with 1.25 μ g of avidin-conjugated alkaline phosphatase (Sigma) per ml for 2 h at room temperature. After a final wash with PBS, IFN- γ or IL-4 spot-forming cells were detected by the addition of 5-bromo-4-chloro-3-indolylphosphate (BCIP)–nitroblue tetrazolium solution (Sigma) and were counted with a stereomicroscope.

In vivo expansion of CTL effectors in tumor protection model. Seven- to 8-week-old female BALB/c mice were immunized i.n. with 10 μ l of vaccine suspension containing 10⁹ CFU on days 0 and 14 as primary and booster immunizations. At day 14 after the last immunization, mice were given subcutaneous injections into the right flank of CT26 target cells which stably express N protein. The tumor sizes were measured with calipers every 3 days. Tumor volumes were calculated according to the formula volume = width² \times length \times 0.52. The data are presented as means \pm standard errors (SE).

Statistical analysis. One-way analysis of variance (ANOVA) and Student's *t* tests were used for comparison of antibody titers, cytotoxicity levels, tumor growth and IFN- γ or IL-4 production by ELISPOT among different groups. All tests were performed using SPSS software. Values of *P* < 0.05 are considered significant.

RESULTS

Construction of *S. enterica* serovar Typhi *pilS::Km^r pilV::N⁺* recombinant strain. The recombinant *S. enterica* serovar Typhi *pilS::Km^r pilV::N⁺* strain carrying the N gene of SARS-CoV was constructed according to the procedures described in Materials and Methods and shown in Fig. 1A. The vaccine strain was confirmed by PCR amplification using the primers outside recombinant sequences (in which the N gene was inserted into the *pilV* gene) (Fig. 1B). The size of the PCR product was 1,500 bp from the *S. enterica* serovar Typhi *pilS::Km^r pilV::N⁺* recombinant vaccine strain (Fig. 1B), while the PCR product

from the *S. enterica* serovar Typhi *pilS::Km^r* parental strain was 300 bp (Fig. 1B). PCR using the primers with insert sequences (Fig. 1C) also indicated that the N gene (1,200 bp) was inserted into the *pilV* gene of the genomic DNA in the *S. enterica* serovar Typhi *pilS::Km^r pilV::N⁺* recombinant strain.

We further examined the N protein expression in the *S. enterica* serovar Typhi *pilS::Km^r pilV::N⁺* recombinant strain by sodium dodecyl sulfate-polyacrylamide gel electrophoresis and Western blot analysis. As shown in Fig. 1D, a protein band of approximately 45 kDa (corresponding to the molecular mass of N protein) in the cell lysates of the *S. enterica* serovar Typhi *pilS::Km^r pilV::N⁺* strain (Fig. 1D, lane 2) was identified but not detected in the cell lysates of the *S. enterica* serovar Typhi *pilS::Km^r* parental strain (Fig. 1D, lane 1). The results demonstrated that the N protein of SARS-CoV was expressed in the newly constructed *S. enterica* serovar Typhi *pilS::Km^r pilV::N⁺* recombinant strain. We have demonstrated the stability of insertion and expression of vaccine strain after several in vitro passages.

***S. enterica* serovar Typhi *pilS::Km^r pilV::N⁺* vaccine strain elicited predominant IgG2a subclass antibody responses.** Serum IgG responses against N protein were determined in the BALB/c mice immunized with the *S. enterica* serovar Typhi *pilS::Km^r pilV::N⁺* recombinant vaccine strain through different immunization routes, including i.n., o.g., i.p., and i.v. Results from ELISA analysis showed that the i.n. immunization route caused the highest levels of anti-N protein-specific IgG and IgG2a (Fig. 2A and B) (i.n. versus o.g., i.p., and i.v.; *P* < 0.05). In addition, both i.n. and o.g. immunization routes induced the highest levels of specific S-IgA among all routes




FIG. 2. Determination of serum antibody responses against SARS-N protein. Each group of BALB/c mice ($n = 6$) was immunized with the *S. enterica* serovar typhi *pilS*:: Km^r *pilV*:: N^+ recombinant strain through the i.n., o.g., i.p., and i.v. routes. The levels (titers) of serum IgG (A), IgG2a (B), and S-IgA (C), produced in immunized mice were determined by ELISA at day 14 after the last immunization. All of the experiments were repeated five times. The data are means \pm SE.

(Fig. 2C). The parental strain (with no insert) and nonimmunized mice (controls) did not induce a detectable antibody response. The data suggested that i.n. immunization was the best route for this recombinant vaccine. Therefore, we chose the i.n. immunization route for the rest of the experiments in this study. In mice, the presence of IgG2a antibodies is indicative of a Th1-biased response since the Th1 cytokines are necessary for this isotype shift in B cells. Our results demonstrated that immunization of mice with the *S. enterica* serovar Typhi *pilS*:: Km^r *pilV*:: N^+ recombinant strain could induce more of an IgG2a Th1-type response than the parental strain. In addition, the recombinant strain could elicit specific mucosal immunity.

***S. enterica* serovar Typhi *pilS*:: Km^r *pilV*:: N^+ recombinant vaccine stimulated specific CTL responses in mice.** Mice were immunized with the recombinant strain or the parental bacterium strain i.n. on day 0 and 14, respectively, and then euthanized at day 28. The specific CTL activities in splenocytes from the immunized mice were examined. Splenocytes from immunized mice were restimulated with the recombinant N protein in vitro as expanded effector cells and incubated with MHC-matched CT26 ($H-2^d$) cells, which stably expressed N protein as target cells. CTL assays showed that the recombinant vaccine strain elicited significantly higher levels of specific CTL responses than the *S. enterica* serovar Typhi *pilS*:: Km^r parental strain and the control group (Fig. 3). We determined that the highest CTL activity was detected when the ratio of effector cells to target cells (E:T) was 2.5:1. These results demonstrated




FIG. 3. Determination of N-specific CTL responses in mice immunized i.n. with the recombinant vaccine strain. Each group of mice ($n = 6$) was immunized i.n. with the *S. enterica* serovar Typhi *pilS*:: Km^r parental strain or *S. enterica* serovar Typhi *pilS*:: Km^r *pilV*:: N^+ recombinant strain on days 0 and 14, respectively. Splenocytes were harvested at day 14 after the last immunization and stimulated by the recombinant N protein as effector cells. MHC-matched CT26 ($H-2^d$) cells, which stably expressed N protein, were used as target cells. All columns showed the mean percentages of special cytotoxicity at an E:T ratio of 2.5:1 in the LDH release assay. The experiments were repeated six times. The data are means \pm SE.




FIG. 4. Cytokine production. Each group of mice ($n = 6$) was immunized with the *S. enterica* serovar Typhi *pilS*::Km^r parental strain or *S. enterica* serovar Typhi *pilS*::Km^r *pilV*::N⁺ recombinant strain. Levels of production of IFN- γ (A) and IL-4 (B) spot-forming-cells (SFC) per 10⁶ splenocytes were assessed by ELISPOT assay using splenocytes isolated from immunized mice. All columns show the mean numbers of SFC; the error bars indicate SD. A significant increase of IFN- γ secretion was observed in mice immunized with the *S. enterica* serovar Typhi *pilS*::Km^r *pilV*::N⁺ recombinant strain. *, $P < 0.05$.

that the *S. enterica* serovar Typhi *pilS*::Km^r *pilV*::N⁺ recombinant vaccine strain could significantly stimulate specific CTL responses in mice.

***S. enterica* serovar Typhi *pilS*::Km^r *pilV*::N⁺ recombinant vaccine induced IFN- γ production.** IFN- γ secretion is indicative of a Th1-biased response, while IL-4 production is indicative of a Th2-biased response. Using the ELISPOT technique, production of IFN- γ and IL-4 was detected from splenocytes in all immunized mouse groups. After stimulation by the recombinant N protein (expressed and purified from *E. coli* BL21), splenocytes from the recombinant strain group induced significantly higher levels of IFN- γ -producing cells (mean \pm SD, 5,270 \pm 244) than those from the parental strain group (mean \pm SD, 2,300 \pm 268) ($P < 0.05$) (Fig. 4A). However, there were no significant differences in the levels of expression of IL-4-producing cells among all groups, including the recombinant vaccine group, parental strain group, and nonimmunized control group (Fig. 4B). This indicated that the recombinant strain induced a Th1-biased immune response.

***S. enterica* serovar Typhi *pilS*::Km^r *pilV*::N⁺ recombinant vaccine inhibited the N-expressing tumors growing in vivo.** Specific cellular immune response is a crucial mechanism of defense against intracellular microorganisms, viruses, and tumors. In this study, BALB/c mice were immunized with the recombinant vaccine or the parental strain. At day 14 after the last immunization, mice were given subcutaneous injections of CT26 target cells stably expressing N protein. The CD8⁺ T cells of BALB/c mice recognized MHC-I-matched CT26 cells and protected BALB/c mice from CT26 tumor challenge. The results showed that the tumor sizes in the recombinant vaccine group were significantly smaller than those in the control parental strain group ($P < 0.05$) (Fig. 5A and B). The sizes of tumors increased significantly in a time-dependent fashion in the parental strain group but insignificantly in the recombinant vaccine group (Fig. 5A). These data showed that the recombinant strain could inhibit the N-expressing tumors growing in vivo by increasing the specific cellular immune response.

DISCUSSION

S. enterica serovar Typhi strains are attractive for their use as vaccine vectors since they can be administered by the natural route of infection (i.e., orally) and are capable of eliciting both systemic and mucosal immune responses (11, 15, 37, 38). Of the more than 2,300 closely related *Salmonella* serovars recognized, *S. enterica* serovar Typhi is the only one that invades the bloodstream and causes systemic infection and immunity exclusively to humans. Some attenuated *S. enterica* serovar Typhi strains (for example, *S. enterica* serovar Typhi Ty21a) have been reported to be used for expression of heterologous antigens or proteins that can be delivered to the immune system in human studies (21). The possibility of using *S. enterica* serovar Typhi as a live vector to express heterologous antigens or deliver them in genetic form in a multivalent vaccine that could protect against several diverse pathogens is an attractive alternative in vaccine development and delivery (4–7, 14, 16, 24).

In this study, the reason we selected the *S. enterica* serovar Typhi *pilS*::Km^r strain as the parental bacterium is that the expression of virulent antigen Vi and the expression of type IVB pili are defective (44). The type IVB pilus of the pathogen has been identified as an important virulence factor that was required for pathogenicity (44). Moreover, the type IVB pilus operon is confined to *S. enterica* serovar Typhi and a few other human-invasive strains such as *Salmonella enterica* serovars Paratyphi C and Dublin (26, 40). The *S. enterica* serovar Typhi mutant (*pilS*::Km^r) strain significantly attenuated the adhesion to and invasion of human intestinal or monocytic cells compared with those of the wild-type strain (28, 44). Of the attenuated *S. enterica* serovar Typhi strains (for example *S. enterica* serovar Typhi Ty21a), the immunogenicity of Ty21a is not remarkable, particularly when the vaccine is administered in capsule form by oral vaccination. Other highly immunogenic *S. enterica* serovar Typhi derivatives made and tested to date have either caused fever or may retain the potential to cause fever in at least some small numbers of volunteer recipients. So the attenuated *S. enterica* serovar Typhi *pilS*::Km^r strain was ex-




FIG. 5. Analysis of specific antitumor activity of CTL responses in immunized mice. (A) Growth curve of tumor cells. *, $P < 0.05$ for the *S. enterica* serovar Typhi *pilS*::Km^r *pilV*::N⁺ group versus *S. enterica* serovar Typhi *pilS*::Km^r parental strain. The data are means \pm SE. (B) Tumor sizes in immunized mice on the 20th day postinjection with 2×10^6 CT26 tumor cells.

amined in this study for its possibility to elicit a full and strong immune response by the natural route of infection. Our results clearly showed that the type IVB pilus operon promoter controlling nucleocapsid gene expression of SARS-CoV in the *S. enterica* serovar Typhi *pilS*::Km^r*pilV*::N⁺ strain could elicit a full specific immune response by i.n. vaccination.

Since plasmid DNA is relatively easily lost, we constructed this attenuated *S. enterica* serovar Typhi vaccine stably expressing the conserved N protein of SARS-CoV by integrating the N gene into the *pilV* gene under the control of *pil* promoter in the chromosome of the strain. This expression of N protein in the vaccine strain is stable after many (more than 20) in vitro passages (data not shown). However, this attenuated *S. enterica* serovar Typhi strain is a Ty2 derivative, EX462, with rifampin-resistant, Vi-negative, β -galactosidase mutant and with a kanamycin cassette inserted to inactive *pilS* (44). Therefore, this strain encodes resistance to two antibiotics, which limits the experimental and clinical studies that can be done with animals. For this reason, future work may be performed to insert other candidate antigen genes (for example, the gene coding for SARS viral spike protein) into the two antibiotic resistance genes (Rif^r and Kam^r) to confer their resistance.

In recent years, the i.n. dosing model has been used to demonstrate the immunogenicity of *S. enterica* serovar Typhi

vaccine strains in a number of other studies (9, 12, 29, 30, 32). For example, *S. enterica* serovar Typhi CVD 908 expressing the major protein gp63 of *Leishmania mexicana mexicana* was shown to be immunogenic through this pathway, eliciting both humoral and cellular immune responses (12). Pickett et al. (32) reported that attenuated *S. enterica* serovar Typhi live-vector vaccine strains were highly immunogenic in mice following i.n. but not o.g. inoculation. Despite the established utility of this murine i.n. model (1, 9), little is known about the immunogenicity of serovar Typhi vaccine strains administered by different routes. We have demonstrated that the i.n. route of immunization caused the highest production of specific IgG, IgG2a, and S-IgA to SARS-CoV N antigens delivered by *S. enterica* serovar Typhi live-vector vaccines compared to the o.g., i.p., and i.v. routes (Fig. 2). In addition, administration of 10^9 CFU of bacteria by the i.v. or i.p. route could cause death of mice (data not shown), while 10^7 CFU of bacteria i.v. or i.p. did not induce enough of an immunogenic response (Fig. 2); therefore, we considered the i.n. route to be the best, although the dose of the i.n. and o.g. application was 100-fold higher than that for the i.p. and i.v., which might have some effects on magnitude of response. More safety concerns about the i.n. route for live bacterial vector strains in humans might be considered for further investigation.

Cell-mediated immune response is a crucial mechanism of defense against most intracellular pathogens and tumors (14, 19), although it remains elusive for the new, emerging SARS-CoV. Our data showed that the recombinant strain could inhibit the N-expressing tumor growing in vivo (Fig. 5) by stimulating specific cellular immune responses. We also observed that the attenuated *S. enterica* serovar Typhi *pilS::Km^r pilV::N⁺* vaccine elicited strong specific CTL responses (Fig. 3) as measured by LDH release assay at an E:T ratio of 2.5/1. There was no significant increase of CTL responses at the E:T ratios of 5/1 and 10/1 (data not shown), and the reasons for these are not known. We speculate that one of the reasons might be due to minimal activity detected following in vitro expansion of memory cells. Another possibility is that CTL activities are not measured by direct MHC-I perforin-mediated lysis experiments.

There are several advantages for the attenuated *S. enterica* serovar Typhi *pilS::Km^r pilV::N⁺* recombinant vaccine. First, it can elicit systemic immunity and it is attenuated (28, 44) and relatively safe. Second, it can be used as a mucosal vaccine vector to deliver pathogen-specific protective epitopes into the mucosa-associated lymphoid tissues. Third, it is easy to manipulate, grows quickly, and costs much less than viral vector vaccines or DNA vector vaccines. Both mucosal and systemic immune responses against the carrier and the foreign antigens may be obtained via the i.n. route (29).

Until now, effective live vaccines that can elicit both mucosal and systemic immune responses against SARS-CoV have been very limited. Our newly constructed attenuated *S. enterica* serovar Typhi *pilS::Km^r pilV::N⁺* vaccine proved to be a highly proficient live-vaccine candidate. Our work also provides insights into how the integration of other foreign genes into the genes under the control of the type IVB pilus promoter in the attenuated *S. enterica* serovar Typhi live vector might be attractive for vaccine delivery and vaccine candidate development against intracellular microorganism infection, respiratory infection, or inhibition of tumor growth. This is due to its ability to stably express foreign antigens and induce mucosal, humoral, and cellular immune responses.

ACKNOWLEDGMENTS

This work was supported by the grants from National Natural Science Foundation of China (no. 30670098, 30470087, and 30570070), the Major State Basic Research Development Program of China ("973" project no. 2005CB522901 and 2006CB504300), the Ministry of Education Scientific Research Foundation for the New Century Outstanding Scholars (NCET-04-0685), Hubei Province Department of Science and Technology (no. 2005S2354, 2005ABC003, 2006ABD007, and 2005AA304B04), and the Hubei Ministry of Public Health (no. JX1B074).

REFERENCES

- Barry, E. M., O. Gomez-Duarte, S. Chatfield, R. Rappuoli, M. Pizza, G. Losonsky, J. Galen, and M. M. Levine. 1996. Expression and immunogenicity of pertussis toxin S1 subunit-tetanus toxin fragment C fusions in *Salmonella typhi* vaccine strain CVD 908. *Infect. Immun.* **64**:4172-4181.
- Buchholz, U. J., A. Bukreyev, L. Yang, E. W. Lamirande, B. R. Murphy, K. Subbara, and P. L. Collins. 2004. Contributions of the structural proteins of severe acute respiratory syndrome coronavirus to protective immunity. *Proc. Natl. Acad. Sci. USA* **101**:9804-9809.
- Bukreyev, A., E. W. Lamirande, U. J. Buchholz, L. N. Vogel, W. R. Elkins, M. St. Claire, B. R. Murphy, K. Subbara, and P. L. Collins. 2004. Mucosal immunization of African green monkeys (*Cercopithecus aethiops*) with an attenuated parainfluenza virus expressing the SARS coronavirus spike protein for the prevention of SARS. *Lancet* **26**:2122-2127.
- Dietrich, G., S. Spreng, I. Gentschev, and W. Goebel. 2000. Bacterial systems for the delivery of eukaryotic antigen expression vectors. *Antisense Nucleic Acid Drug Dev.* **10**:391-399.
- Dietrich, G., I. Gentschev, J. Hess, J. B. Ulmer, S. H. Kaufmann, and W. Goebel. 1999. Delivery of DNA vaccines by attenuated intracellular bacteria. *Immunol. Today* **20**:251-253.
- Dietrich, G., and W. Goebel. 2000. DNA vaccine delivery by attenuated intracellular bacteria. *Subcell. Biochem.* **33**:541-557.
- Drabner, B., and C. A. Guzman. 2001. Elicitation of predictable immune responses by using live bacterial vectors. *Biomol. Eng.* **17**:75-82.
- Eisenstein, T. K. 1999. Mucosal immune defense: the *Salmonella typhimurium* model, p. 51-109. *In* Y. Paterson (ed.), *Intracellular bacterial vaccines*. Wiley-Liss, Inc., New York, NY.
- Galen, J. E., O. G. Gomez-Duarte, G. A. Losonsky, J. L. Halpern, C. S. Lauderbaugh, S. Kaintuck, M. K. Reymann, and M. M. Levine. 1997. A murine model of intranasal immunization to assess the immunogenicity of attenuated *Salmonella typhi* live vector vaccines in stimulating serum antibody responses to expressed foreign antigens. *Vaccine* **15**:700-708.
- Garmory, H. S., K. A. Brown, and R. W. Titball. 2002. *Salmonella* vaccines for use in humans: present and future perspectives. *FEMS Microbiol. Rev.* **26**:339-353.
- Gonzalez, C., D. Hone, F. R. Noriega, C. O. Tacket, J. R. Davis, G. Losonsky, J. P. Nataro, S. Hoffman, A. Malik, and E. Nardin. 1994. *Salmonella typhi* vaccine strain CVD 908 expressing the circumsporozoite protein of *Plasmodium falciparum*: strain construction and safety and immunogenicity in humans. *J. Infect. Dis.* **69**:927-931.
- Gonzalez, C. R., F. R. Noriega, S. Huerta, A. Santiago, M. Vega, J. Paniagua, V. Ortiz-Navarrete, A. Isibasi, and M. M. Levine. 1998. Immunogenicity of a *Salmonella typhi* CVD 908 candidate vaccine strain expressing the major surface protein gp63 of *Leishmania mexicana mexicana*. *Vaccine* **16**:1043-1052.
- He, Y., J. Li, S. Heck, S. Lustigman, and S. Jiang. 2006. Antigenic and immunogenic characterization of recombinant baculovirus-expressed severe acute respiratory syndrome coronavirus spike protein: implication for vaccine design. *J. Virol.* **80**:5757-5767.
- Hess, J., U. Schaible, B. Raupach, and S. H. Kaufmann. 2000. Exploiting the immune system: toward new vaccines against intracellular bacteria. *Adv. Immunol.* **75**:1-88.
- Hohmann, E. L., C. A. Oletta, K. P. Killeen, and S. I. Miller. 1996. *phoP*/*phoQ*-deleted *Salmonella typhi* (Ty800) is a safe and immunogenic single dose typhoid fever vaccine in volunteers. *J. Infect. Dis.* **173**:1408-1414.
- Hone, D. M., S. R. Attridge, B. Forrest, R. Morona, D. Daniels, J. T. LaBrooy, R. C. Bartholomeusz, D. J. C. Shearman, and J. Hackett. 1988. A *galE* via (Vi antigen-negative) mutant of *Salmonella typhi* Ty2 retains virulence in humans. *Infect. Immun.* **56**:1326-1333.
- Ishii, K., H. Hasegawa, N. Nagata, T. Mizutani, S. Morikawa, T. Suzuki, F. Taguchi, M. Tashiro, T. Takemori, T. Miyamura, and Y. Tsunetsugu-Yokota. 2006. Induction of protective immunity against severe acute respiratory syndrome coronavirus (SARS-CoV) infection using highly attenuated recombinant vaccinia virus DIs. *Virology* **351**:368-380.
- Jin, H., C. Xiao, Z. Chen, Y. Kang, Y. Ma, K. Zhu, Q. Xie, Y. Tu, Y. Yu, and B. Wang. 2005. Induction of Th1 type response by DNA vaccinations with N, M, and E genes against SARS-CoV in mice. *Biochem. Biophys. Res. Commun.* **328**:979-986.
- Kaufmann, S. H. E. 1998. Immunity to intracellular bacteria, p. 1335-1371. *In* W. E. Paul (ed.), *Fundamental immunology*. Lippincott-Raven, New York, NY.
- Kim, T. W., J. H. Lee, C.-F. Hung, S. Peng, R. Roden, M.-C. Wang, R. Viscidi, Y.-C. Tsai, L. He, P.-J. Chen, D. A. K. Boyd, and T.-C. Wu. 2004. Generation and characterization of DNA vaccines targeting the nucleocapsid protein of severe acute respiratory syndrome coronavirus. *J. Virol.* **78**:4638-4645.
- Kotton, C. N., and E. L. Hohmann. 2004. Enteric pathogens as vaccine vector for foreign antigen delivery. *Infect. Immun.* **72**:5535-5547.
- Lee, J. S., K. S. Shin, J. G. Pan, and C. J. Kim. 2000. Surface-displayed viral antigens on *Salmonella* carrier vaccine. *Nat. Biotechnol.* **18**:645-648.
- Lee, J.-S., H. Poo, D. P. Han, S.-P. Hong, K. Kim, M. W. Cho, E. Kim, M.-H. Sung, and C.-J. Kim. 2006. Mucosal immunization with surface-displayed severe acute respiratory syndrome coronavirus spike protein on *Lactobacillus casei* induces neutralizing antibodies in mice. *J. Virol.* **80**:4079-4087.
- Levine, M. M., J. E. Galen, E. Barry, F. Noriega, C. Tacket, M. B. Szein, S. Chatfield, G. Dougan, G. Losonsky, and K. Kotloff. 1997. Attenuated *Salmonella typhi* and *Shigella* as live oral vaccines and as live vectors. *Behring Inst. Mitt.* **98**:120-123.
- Marra, M. A., S. J. M. Jones, C. R. Astell, R. A. Holt, A. Brooks-Wilson, Y. S. N. Butterfield, et al. 2003. The genome sequence of the SARS-associated coronavirus. *Science* **300**:1399-1404.
- Morris, C., C. K. Tam, T. S. Wallis, P. W. Jones, and J. Hackett. 2003. *Salmonella enterica* serovar Dublin strains which are Vi antigen-positive use type IVB pili for bacterial self-association and human intestinal cell entry. *Microb. Pathog.* **35**:279-284.
- Nardelli-Haeffiger, D., J. Benyacoub, R. Lemoine, S. Hopkins-Donaldson, A.

- Potts, F. Hartman, J. P. Kraehenbuhl, and P. Grandi. 2001. Nasal vaccination with attenuated *Salmonella typhimurium* strains expressing the hepatitis B virus nucleocapsid: dose response analysis. *Vaccine* **19**:2854–2861.
28. Pan, Q., X.-L. Zhang, H.-Y. Wu, P.-W. He, F. Wang, M.-S. Zhang, J.-M. Hu, B. Xia, and J. Wu. 2005. Aptamers that preferentially bind the type IVB pili and inhibit human monocytic-cell invasion by *Salmonella enterica* serovar Typhi. *Antimicrob. Agents Chemother.* **49**:4052–4060.
 29. Pasetti, M. F., R. J. Anderson, F. R. Noriega, M. M. Levine, and M. B. Szein. 1999. Attenuated Δ guaBA *Salmonella typhi* vaccine strain CVD915 as a live vector utilizing prokaryotic or eukaryotic expression systems to deliver foreign antigens and elicit immune responses. *Clin. Immunol.* **92**:76–89.
 30. Pasetti, M. F., T. E. Pickett, M. M. Levine, and M. B. Szein. 2000. A comparison of immunogenicity and *in vivo* distribution of *Salmonella enterica* serovar Typhi and Typhimurium live vector vaccines delivered by mucosal routes in the murine model. *Vaccine* **18**:3208–3213.
 31. Peng, H., L. T. Yang, L. Y. Wang, J. Li, J. Huang, Z. Q. Lu, R. A. Koup, R. T. Bailer, and C. Y. Wu. 2006. Long-lived memory T lymphocyte responses against SARS coronavirus nucleocapsid protein in SARS-recovered patients. *Virology* **351**:466–475.
 32. Pickett, T. E., M. F. Pasetti, J. E. Galen, M. B. Szein, and M. M. Levine. 2000. *In vivo* characterization of the murine intranasal model for assessing the immunogenicity of attenuated *Salmonella enterica* serovar Typhi strains as live mucosal vaccines and as live vectors. *Infect. Immun.* **68**:205–213.
 33. Saif, L. J. 2004. Animal coronavirus vaccines: lessons for SARS. *Dev. Biol.* **119**:129–140.
 34. See, R. H., A. N. Zakhartchouk, M. Petric, D. J. Lawrence, C. P. Mok, R. J. Hogan, et al. 2006. Comparative evaluation of two severe acute respiratory syndrome (SARS) vaccine candidates in mice challenged with SARS coronavirus. *J. Gen. Virol.* **87**:641–650.
 35. Spruth, M., O. Kistner, H. Savidis-Dacho, E. Hitter, B. Crowe, M. Gerencer, P. Bruhl, L. Grillberger, M. Reiter, C. Tauer, W. Mundt, and P. N. Barrett. 2006. A double-inactivated whole virus candidate SARS coronavirus vaccine stimulates neutralizing and protective antibody responses. *Vaccine* **24**:652–659.
 36. Surjitt, M., B. Liu, P. Kumar, V. T. Chow, and S. K. Lal. 2004. The nucleocapsid protein of the SARS coronavirus is capable of self-association through a C-terminal 209 amino acid interaction domain. *Biochem. Biophys. Res. Commun.* **317**:1030–1036.
 37. Szein, M. B., and G. Mitchell. 1997. Recent advances in immunology: impact on vaccine development, p. 99–125. *In* M. M. Levine, G. C. Woodrow, J. B. Kaper, and G. S. Cobon (ed.), *New generation vaccines*. Marcel Dekker, Inc., New York, NY.
 38. Tacket, C. O., M. B. Szein, G. A. Losonsky, S. S. Wasserman, J. P. Natario, R. Edelman, D. Pickard, G. Dougan, S. N. Chatfield, and M. M. Levine. 1997. Safety of live oral *Salmonella typhi* vaccine strains with deletions in *htrA* and *aroC aroD* and immune response in humans. *Infect. Immun.* **65**:452–456.
 39. Takeda, J., Y. Sato, H. Kiyosawa, T. Mori, S. Yokoya, A. Irisawa, M. Miyata, K. Obara, T. Fujita, T. Suzuki, R. Kasukawa, and A. Wanaka. 2000. Anti-tumor immunity against CT26 colon tumor in mice immunized with plasmid DNA encoding β -galactosidase fused to an envelope protein of endogenous retrovirus. *Cell. Immunol.* **204**:11–18.
 40. Tam, C. K. P., J. Hackett, and C. Morris. 2004. *Salmonella enterica* serovar Paratyphi C carries an inactive shufflon. *Infect. Immun.* **72**:22–28.
 41. Wang, J., J. Wen, J. Li, J. Yin, Q. Zhu, H. Wang, et al. 2003. Assessment of immunoreactive synthetic peptide from the structural proteins of severe acute respiratory syndrome coronavirus. *Clin. Chem.* **49**:1989–1996.
 42. Yang, L. T., H. Peng, Z. L. Zhu, G. Li, Z. T. Huang, Z. X. Zhao, R. A. Koup, R. T. Bailer, and C. Y. Wu. 2006. Long-lived effector/central memory T-cell responses to severe acute respiratory syndrome coronavirus (SARS-CoV) S antigen in recovered SARS patients. *Clin. Immunol.* **120**:171–178.
 43. Reference deleted.
 44. Zhang, X.-L., I. S. M. Tsui, C. M. C. Yip, A.W. Y. Fung, D. K.-H. Wong, X. Dai, Y. Yang, J. Hackett, and C. Morris. 2000. *Salmonella enterica* serovar Typhi uses type IVB pili to enter human intestinal epithelial cells. *Infect. Immun.* **68**:3067–3073.
 45. Zhang, X. L., C. Morris, and J. Hackett. 1997. Molecular cloning, nucleotide sequence, and function of a new site-specific recombinase encoded in the major “pathogenicity island” of *Salmonella* Typhi. *Gene* **202**:139–146.
 46. Zhu, Y., M. Liu, W. Zhao, J. Zhang, X. Zhang, K. Wang, C. Gu, Y. Li, C. Zheng, G. Xiao, H. Yan, J. Zhang, D. Guo, P. Tien, and J. Wu. 2005. Isolation of virus from a SARS patient and genome-wide analysis of genetic mutations related to pathogenesis and epidemiology from 47 SARS-CoV isolates. *Virus Genes* **30**:93–102.